

TCCC Journal Watch

February 2010

Abernathy J, Reeves S: Airway catastrophes. Curr Opin Anaesthesiol 2010;23(1):41-46

Belmont P, Goodman G, Zacchilli M, et al: Incidence and epidemiology of combat injuries sustained during “the surge” portion of operation Iraqi Freedom by a U.S. Army brigade combat team. J Trauma 2010;68(1):204-210

Bouillon B, Brohi K, Hess J, et al: Educational initiative on critical bleeding in trauma: Chicago, July 11-13, 2008. J Trauma 2010;68(1):225-230

Burgert J: Intraosseous infusion of blood products and epinephrine in an adult patient in hemorrhagic shock. AANA J 2009;77(5):359-363

Dong W, Cai B, Peña G, et al: Ethyl pyruvate prevents inflammatory responses and organ damage during in porcine hemorrhage. Shock 2009;Epub ahead of print

Eastridge B, Costanzo G, Jenkins D, et al: Impact of joint theater trauma system initiatives on battlefield injury outcomes. Am J Surg 2010;198(6): 852-857

Flint N, Russell W, Thompson J: Comparison of different methods of ventilation via cannula cricothyroidotomy in a trachea-lung model. Br J Anaesth 2009;103(6):891-895

Franschman G, Greuters S, Loer S, et al: Prehospital treatment guidelines in severe traumatic brain injury: What really happens outside the hospital? Resuscitation 2009;Epub ahead of print.

Goon S, Stephens R, Smith H: The emergency airway. Br J Hosp Med 2010;70(12):M186-188

Hargreaves S, Lacasia C, Groom P: Regional audit- availability of equipment and training in emergency cricothyroidotomy. Anaesthesia 2010;65(1):107

Haut E, Kalish B, Efron D, et al: Spine immobilization in penetrating trauma: more harm than good? J Trauma 2010;68(1):115-121

Hetzler M, Rick G: Damage control resuscitation for the special forces medic- simplifying and improving prolonged trauma care-part two. J Spec Oper Med 2009;9(4):53-62

Holcomb J, Spinella P: Optimal use of blood in trauma patients. Biologicals 2010; Epub ahead of print

Hoyt B: Blood and war-lest we forget. J Am Coll Surg 2009;209(6):681-6

Jiao W, Kiang J, Cary L, et al: COX-2 inhibitors are contraindicated for treatment of combined injury. Radiat Res 2009;172(6):686-697

Lamhaut L, Dagrón C, Apriotesei R, et al: Comparison of intravenous and intraosseous access by pre-hospital medical emergency personnel with and without CBRN protective equipment. Resuscitation 2009; Epub ahead of print

Lee J, Kim K, Jo Y, et al: Early norepinephrine infusion delays cardiac arrest after hemorrhagic shock in rats. J Emerg Med 2009;37(4):376-382

Lew T, Walker J, Wenke J, et al: Characterization of craniomaxillofacial battle injuries sustained by the United States service members in the current conflicts of Iraq and Afghanistan. J Oral Maxillofac Surg 2010;68(1): 3-7

Ling G, Rhee P, Ecklund J: Surgical innovations arising from the Iraq and Afghanistan wars. Annu Rev Med 2010;61:457-468

Morrison J, Mountain A, Galbraith K, et al: Penetrating pelvic battlefield trauma; internal use of chitosan-based haemostatic dressings. Injury 2010;41(2):239-241

Napolitano L, Kurek S, Luchette F, et al: Clinical practice guideline: red blood cell transfusion in adult trauma and critical care. Crit Care Med 2009;37(12):3124-3157

Nelson B, Melnick E, Li J: Portable ultrasound for remote environments, part i: feasibility of field deployment. J Emerg Med 2010;Epub ahead of print

Rockswold G, Solid C, Paredes-Andrade E, et al; Hypertonic saline and its effect on intracranial pressure, cerebral perfusion pressure, and brain tissue oxygen Neurosurgery 2009;65(6):1035-1041

Sánchez-Morillas L, Rojas Pérez-Ezquerria P, Reaño-Martos M, et al: Systemic anaphylaxis caused by moxifloxacin. Allergol Immunopathol 2010;Epub ahead of print

Schuster K, Lofthouse R, Moore C, et al: Pulseless electrical activity, focused abdominal sonography for trauma, and cardiac contractile activity as predictors of survival after trauma. J Trauma 2009;67(6):1154-1157

Sen A, Gulati A: Use of Magnesium in traumatic brain injury. Neurotherapeutics 2010;7(1):91-99

Sharp C, Mangram A, Lorenzo M, et al: A major metropolitan “field amputation” team: a call to arms ... and legs. J Trauma 2009;67(6): 1158-1161

Sondén A, Rocksén D, Riddez L, et al: Trauma attenuating backing improves protection against behind armor blunt trauma. J Trauma 2009;67 (6):1191-1199

Ward K, Archambault R, Mersfelder T: Severe adverse skin reaction to nonsteroidal anti-inflammatory drugs: a review of literature. Am J Health Syst Pharm 2010;67(3):206-213

November 2009

Alam H, Bice L, Butt M, et al: Testing of blood products in a polytrauma model: results of a multi-institutional randomized preclinical trial. J Trauma 2009;67(4):856-864

Allard C, Scarpelini S, Rhind S, Baker A, et al: Abnormal coagulation test are associated with progression of traumatic intracranial hemorrhage. J Trauma 2009;67(5):959-967

Ambrus R, Kocbek P, Kristi J, et al: Investigation of preoperation parameters to improve the dissolution of poorly water-soluble meloxicam. Int J Pharm 2009;381(2):153-159

Arnaud F, Teranishi K, Tomori T, et al: Comparison of 10 hemostatic dressing in a groin puncture model in swine. J Vasc Surg 2009;50(3):632-639

Batista M, Bravin A, Lopes L, et al: Pressor response to fluid resuscitation in endotoxic shock: involvement of vasopressin. Crit Care Med 2009;37(11):2968-2972

Blackwell T, Kline J, Willis J, et al: Lack of association between prehospital response times and patient outcomes. Prehosp Emerg Care 2009;13:444-450

Buckenmaier C, Rupprecht C, McKnight G, et al: Pain following battlefield injury and evacuation: a survey of 110 casualties for the wars in Iraq and Afghanistan. Pain Med 2009;Oct: Epub ahead of print

Cabrales P: Low dose nitrite enhances perfusion after fluid resuscitation from hemorrhagic shock. Resuscitation 2009;Oct:Epub ahead of print

Cheatham M, Safcsak K: Is the evolving management of intra-abdominal hypertension and abdominal compartment syndrome improving survival? Crit Care Med 2009;Sept;Epub ahead of print

Cho R, Bakken H, Reynolds M, et al: Concomitant cranial and ocular combat injuries during Operations Iraq Freedom. J Trauma 2009;67(3) 5126-520

Chung K, Wolf S, Cancio L, et al: Resuscitation of severely burned military casualties; fluid begets more fluid. J Trauma 2009;67(2):231-237

Cobas M, De la Peña M, Manning R et al: Prehospital intubations and mortality: a level 1 trauma center perspective. Anesth Analg 2009; 109(2): 489-493

Cotton B, Jerome R, Collier B, et al: Guidelines for prehospital fluid resuscitation in the injured patient. J Trauma 2009;67(2):389-402

Elliot M, Jallo J, Barbe M, et al: Hypertonic saline attenuates tissue loss and astrocyte hypertrophy in a model of traumatic brain injury. Brain Res 2009:Epub ahead of print

Fatovich D, Jacobs I: The relationship between remoteness and trauma deaths in Western Australia. J Trauma 2009;67(5):910-914

Flint N, Russell W, Thompson J: Comparison of different methods of ventilation via cannula cricothyroidotomy in a trachea-lung model. Br J Anaesth 2009;Sept:Epub ahead of print.

Fludger S, Bell A: Tourniquet application in a rural Queensland HEMS environment. Air Med J 2009;28(6):291-293

Franschman G, Peerdeman S, Greuters S, et al: Prehospital endotracheal intubation in patients with severe traumatic brain injury: guidelines verses reality. Resuscitation 2009;80(10):1147-1151

Gerhardt R, Matthews J, Sullivan S: The effect of systemic antibiotic prophylaxis and wound irrigation on penetrating combat wounds in a return-to-duty population. Prehosp Emerg Care 2009;13(4):500-504

Gonzalez C, Brunstein A: Training for emergencies. J Trauma 2009;67(2):S100-105

Greenburg A: The ideal blood substitute. Crit Care 2009;25:415-424

Hardaway R: One surgeon's army experience with "wound shock" from Pearl Harbor to present. Mil Med 2009;174(9):944-947

Hecht-Dolnick M, Barkan H, Taharka A, et al: Hetastarch increases the risk of bleeding complications in patients after off-pump coronary bypass surgery: a randomized clinical trial. J Thorac Cardiovasc Surg 2009;138(3): 703-711

Holmes J, Wisner D, McGahan J, et al: Clinical prediction rules for identifying adults at very low risk for intra-abdominal injuries after blunt trauma. Annals of Emerg Med;54(4):575-584

Hospital del Henares, Madrid, Spain: Steroid use in acute lung injury. Crit Care Med 2009;37(9):2680

Johansson P, Stissing T, Bochsén L, et al: Thrombelastography in assessing coagulopathy in trauma. Scan J Trauma Resusc Emerg Med 2009;17(1):45

Kaspar R, Griffith M, Mann P, et al: Association of bacterial colonization at the time of presentation to a combat support hospital in a combat zone with subsequent 30-day colonization or infection. Mil Med 2009;174:899-903

Kattail D, Furlan J, Fehlings M: Epidemiology and clinical outcomes of acute spine trauma and spinal cord injury: experience from a specialized spine trauma center in Canada in comparison with a large national registry. J Trauma 2009;67(5):936-943

Kheirabadi B, Scherer M, Estep J, et al: Determination of efficacy of new hemostatic dressings in a model of extremity arterial hemorrhage in swine. J Trauma 2009;67(3):459-460

Kiraly L, Underwood S, Differding J, et al: Transfusion of aged packed red blood cells results in decreased tissue oxygenation in critically injured trauma patient. J Trauma 2009;67(1):29-32

Knapp B, Tsuchitani S, Sheele J, et al: Prospective evaluation of an emergency medical services-administered alternative transport protocol. Prehosp Emerg Care 2009;13:432-436

Leidel B, Kirchhoff C, Bogner V, et al: Is there intraosseous access route fast and efficacious compared to conventional central venous catheterization in adult patients under resuscitation in the emergency department? A prospective study observational pilot study. Patient Saf Surgery 2009;8:24

Levitan R, Bortle C, Snyder T, et al: Use of a battery operated needle driver for intraosseous access by novice users: skill acquisition with cadavers. Annals of Emerg Med 2009;54(5): 692-694

Mahambrey T, Fowler A, Pinto R, et al: Early massive transfusion in trauma patients: Canadian single-centre retrospective cohort study. Can J Anaesth 2009;56(10):740-750

McQuay N, Cipolla J, Franges E, et al: The use of recombinant activated factor VIIa in coagulopathic traumatic brain injury requiring emergent craniotomy: is it beneficial? J Neurosurg 2009;111(4):666-671

Menon D, Zahed C: Prediction of outcome in severe traumatic brain injury. Curr Opin Crit Care 2009;15(5):437-441

Metzger J, Eastman A, Benitez, et al: The livesaving potential of specialized on-scene medical support for urban tactical operations. Prehosp Emerg Care 2009;13(4):528-531

Mistry N, Bleetman A, Roberts K: Chest decompression during the resuscitation of patients in prehospital traumatic cardiac arrest. Emerg Med J 2009;26(10):738-740

Nishijima D, Zehtabchi S: The efficacy of recombinant activated Factor VIIa in severe trauma. Annals of Emerg Med 2009;54(5):737-744

Parker P: Emergency Tourniquet use. Ann Surg 2009;250(3):496-497

Paxton J, Knuth T, Klausnert H; Proximal humerus intraosseous infusion: a preferred emergency venous access. J Trauma 2009;67(3):606-611

Ragal B, Allred C, Brevard S, et al: Fracture of the thoracolumbar spine sustained by soliders in vehicles attacked by improvised explosive devices. Spine 2009: Epub ahead of print.

Sarkar D, Demma A, Stulz D, et al: Expect the unexpected: two cases of penetrating head and neck trauma for Operation Iraqi Freedom. Ear Nose Throat J 2009;88(9):E19-21

Shavit I, Hoffmann Y, Galbraith R, et al: Comparison of two mechanical intraosseous infusion devices: a pilot, randomized crossover trial. Resuscitation 2009;80(9):1029-1033

Solomkin J, Zhao Y, Ma E, et al: Moxifloxacin is non-inferior to combination therapy with ceftriaxone plus metronidazole in patients with community-origin complicated intra-abdominal infections. Int J Antimicrob Agents 2009: Epub ahead of print

Stansbury L, Dutton R, Stein D, et al: Controversy in trauma resuscitation: do ratios of plasma to red blood cells matter? Transfusion Med Rev 2009;23(4):255-265

Sztajnkrzyer M: Needle thoracostomy by non-medical law enforcement personnel: preliminary data on knowledge retention. Prehosp Disaster Med 2009;23(6):553-557

Thomas R, McManus J, Johnson A, et al: Ocular injury reduction for ocular protection use in current combat operations. J Trauma 2009;66:S99-1032

Towner K: Acinetobacter: an old friend, but a new enemy. J Hosp Infect 2009;Epub ahead of print

Vick-Fragoso R, Hernández-Oliva G, Cruz-Alcázar J, et al: Efficacy and safety sequential intravenous/oral Moxifloxican vs intravenous/oral Amoxicillin/Clavulanate for complicated skin and skin structure infections. Infections;Epub ahead of print.

Von Elm E, Schoettker P, Henzi I, et al: Pre-hospital tracheal intubation in patients with traumatic brain injury; systematic review of current evidence. Br J Anaesth 2009;103(3):371-386

Wafaisade A, Lefering R, Tjardes T, et al: Acute coagulopathy in isolated blunt traumatic brain injury. Neurocrit Care 2009;Epub ahead of print.

Yaghoubian A, Lewis R, Putnam B, et al: Reanalysis of prehospital intravenous fluid administration in patients with penetrating truncal injury and field hypertension. Am Surg 2009;73(10):1027-1030

Zhang R, Hunter R, Gonzalez E, et al: Poloxamer 188 prolongs survival of hypotensive resuscitation and decreases vital tissue injury after full resuscitation. Shock 2009;32(4):442-450

August 2009

Allison C, Trunkey D: Battlefield trauma, traumatic shock and consequences: war-related advances in critical care. Crit Care Clin 2009;25:31-45

Beilman G, Blondet j, Nelson T, et al: Early hypothermia in severely injured trauma patients is a significant risk factor for multiple organ dysfunction syndrome but not mortality. Ann Surgery 2009;249:845-850

Brenner T, Gries A, Helm M, et al: Intraosseous infusion systems in the prehospital setting. Resuscitation 2009;80:607

Brown M, Daya M, Worley J, et al: Experience with chitosan dressing in a civilian EMS system.

Cai B, Deitch E, Grande D, et al: Anti-inflammatory resuscitation improves survival in hemorrhage with trauma. J Trauma 2009;66:1632-1640

Carter A, Davis K, Evans L, et al: Information loss in emergency medical services handover trauma patients. Prehosp Emerg Care 2009;13:280-285

Champion H, Holcomb J, Young L: Injuries from explosions: physics, biophysics, pathology, and required research focus. J Trauma 2009;66:1468-1477

Chen L, Resiner A, Gribok A, et al: Exploration of prehospital vital sign trends for the prediction of trauma outcomes. Prehosp Emerg Care 2009;13:286-294

Cobas M, De la Peña M, Manning R, et al: Prehospital intubations and mortality: a level 1 trauma center perspective. Anesth Analg 2009;109:489-493

Cooper E, Bateman S, Muir W: Evaluation of hyperviscous fluid resuscitation in a canine model of hemorrhagic shock: a randomized, controlled study.

Denver Metro Airway Study Group: A prospective multicenter evaluation of prehospital airway management performance in a large metropolitan region. Prehosp Emerg Care 2009;13:304-310

Dougherty A, Mohrie C, Galarneau M, et al: Battlefield extremity injuries in Operation Iraq Freedom. Injury 2009;40:772-777

Ford N, Mills E, Zachariah R, et al: Ethics of conducting research in conflict settings. Confl Health 2009;10:3-7

Gauna AA, Kang SK, Triano ML, et al: Oral Transmucosal fentanyl citrate for dyspnea in terminally ill patients: an observational case series. J Palliat Med 2009;11:643-648

Gerritse B, Scheffer G, Draaisma J: Prehospital intraosseous access with the bone injection gun by a helicopter-transported emergency medical team. J Trauma 2009;66:1739-1741

Huang C, Kao K, Hsu K, et al: Effects of hydroxyethyl starch resuscitation on extravascular lung water and pulmonary permeability in sepsis-related acute respiratory distress syndrome. Crit Care Med 2009;37:1948-1955

Johannigman J, Branson R, Edwards M: Closed loop control of inspired oxygen concentration in trauma patients. J Amer Col of Surgeon 2009;208:763-768

Kilbourne M: Hemostatic efficacy of modified amylopectin powder in a lethal porcine model of extremity arterial injury. Ann Emerg Med 2009;53:804-810

Kucewicz J, Bailey M, Kaczkowski P, et al: Focused ultrasound: concept for automated transcutaneous control of hemorrhage in austere settings. Aviat Space Environ Med;2009;80:391-394

Lam N, Hagberg C, Bassili L: Use of the video laryngoscopy for combitube exchange in a difficult airway. J Clin Anesth 2009;21:294-296

Langeron O, Birenbaum A, Amour J: Airway management in trauma. Minerva Anestesiol 2009;75:307-311

Levitan R, Bortle C, Snyder T, et al: Use of a battery-operated needle driver for intraosseous access by novice users: skill acquisition with cadavers. Ann Emerg Med 2009;Epub ahead of print.

Luck R, Haines C, Mull C: Intraosseous access. J Emerg Med 2009;Epub ahead of print

Lundgren J, Henriksson O, Pretorius T, et al: Field torso-warming modalities: a comparative study using a human model. Prehops Emerg Care 2009;13:371-378

Manring M, Hawk A, Calhoun J, et al: Treatment of war wounds: a historical review. Clin Orthop Relat Res 2009;Epub ahead of print

Martínez-Odrizola P, Gutiérrez-Macías A, Ibarria-Lahuerta J, et al: Meloxicam as a cause of drug-induced autoimmune hepatitis. Dig Dis Sci 2009;Epub of print

McArdle G, McAuley D, McKinley A, et al: Preliminary results of a prospective randomized trial of restrictive versus standard fluid regime in elective open abdominal aortic aneurysm repair. Ann of Surgery 2009;Epub ahead of print.

McQuay N, Cipolla J, Franges E, et al: The use of recombinant activated factor VIIa in coagulopathic traumatic brain injuries requiring emergent craniotomy: is it beneficial? J Neurosurg 2009;Epub ahead of print

Mullins J, Harrahill M: Use of a tourniquet after a gunshot wound to the thigh. J Emerg Nurs 2009;35:265-267

Neparidze N, Kriz N, Ornstein D: Hetastarch haemophilia . Haemophilia 2009;15:976-979

Otero V, Posse J, Carmona I, et al: Efficacy of fluoroquinolones against pathogenic oral bacteria. Mini Rev Med Chem 2009;Epub ahead of print.

Plurad D, Chandrasoma S, Best C, et al: A complication of intracorporeal use of quickclot for pelvic hemorrhage. J Trauma 2009;66:1482-1484

Ramasamy A, Harrison S, Lasrado I, et al: A review of casualties during the Iraq insurgency 2006- a British field hospital experience. Injury 2009;40:493-497

Ramasamy A, Midwinter M, Mahoney P, et al: Learning the lessons from conflict: Pre-hospital cervical spine stabilization following ballistic neck trauma. Injury 2009;Epub ahead of print

Sambasivan C, Cho S, Zink K, et al: A highly porous silica and chitosan-based hemostatic dressing is superior in controlling hemorrhage in a severe groin injury model in swine. Am J Surgery 2009;197:576-580

Schmid M, Mang H, Ey K, et al: Prehospital airway management on rescue helicopters in the United Kingdom. Anaesthesia 2009;64:625-631

Streets C: Lessons from the battlefield in the management of major trauma. Br J Surg 2009;96:831-832

Tam R, Maloney J, Gaboury I, et al: Review of endotracheal intubations by Ottawa Advanced Care Paramedics in Canada. Prehosp Emerg Care 2009;13:311-315

Thaveau F, Zoll J, Bouitbir J, et al: Contralateral leg as a control during skeletal muscle ischemia-reperfusion. J Surg Res 2009;155:65-69

Wang H, Weaver M, Abo B, et al: Ambulance stretcher adverse events. Qual Saf Health Care 2009;18:213-216

April 2009

Archan S, Prause G, Gumpert R, Seibert FJ, Kúgler B: Cricothyroidotomy on the scene in a patient with severe facial trauma and difficult neck anatomy. AM J Emerg Med 2009;27:133

Bauman RA, Ling GS, Tong L, et al: An introductory characterization of a combat casualty care-relevant swine model of closed head injury resulting from exposure to explosive blast. J Neurotrauma 2009 Feb 12; Epub ahead of print

Black IH, McManus J: Pain Management in current combat operations. Prehosp Emerg Care 2009;13:223-227

Bledsoe B, Gandy W: The Disappearing Endotracheal Tube: Historic skill threatened by lack of practice and new devices. JEMS 2009;34:88-99

Bruns B, Gentilello L, Elliot A, Shafi S: Prehospital hypotension redefined. J Trauma 2008;65:1217-1221

Bullard-Berent J: Intraosseous infusion: an instructional program for healthcare providers, revised edition (CD-ROM) Prehosp Emerg Care 2009;13:266

Cai B, Chen F, Lin X, et al: Anti-inflammatory adjuvant in resuscitation fluids improves survival in hemorrhage: Crit Care Med 2009;37:1133-1134

Chan SS: Tension pneumothorax managed without immediate needle decompression. J Emerg Med 2009;36:242-245

David JS, Dubien PY, Capel O, Peguet O, Gueugniaud PY: Intraosseous infusion using the bone injection gun in the prehospital setting. Resuscitation 2009;80:384-385

Devlin JJ, Kircher S, Kozen BG, Littlejohn LF, Johnson AS: Comparison of ChitoFlex®, CELOX, and QuikClot® in Control Hemorrhage. J Emerg Med 2009; Apr 1 - Epub ahead of print

Fatovich DM, Jacobs IG: The relationship between remoteness and trauma deaths in Western Australia. J Trauma 2008 Aug 15 - Epub ahead of print

Gehardt RT, De Lorenzo RA, Oliver J, Holcomb JB, Pfaff JA: Out-of-hospital combat casualty care in the current war in Iraq. Ann Emerg Med 2009;53:169-172

Hartog C, Brunkhorst F, Reinhart K: Hydroxyethyl starch 130/0.4 and surgical blood loss. Anesth Analg 2009;108:672

Hasan SM, Shoaib MH, Hassan F, Rehman I: Bioequivalence studies of two brands of meloxicam tablets in healthy Pakistani volunteers. Pak J Pharm Sci 2009;22:199-204

Hassis VJ, Schino MA, Tepas JJ, et al: Definitive establishment of airway control is critical for optimal outcome in lower cervical spinal cord injury. J Trauma 2008;65:1328-1332

Holcomb JB, McManus JG, Kerr ST, Pusateri AE: Needle versus tube thoracostomy in a swine model of traumatic tension hemopneumothorax Prehosp Emerg Care. 2009;13:18-27

Hurtado TR, Montoya C: Trauma scissors vs. the rescue hook, exposing a simulated patient: a pilot study. J Emerg Med 2009;36:232-235

Johannigman JA, Branson R, Lecroy D, Beck G: Autonomous control of inspired oxygen concentration during mechanical ventilation of the critically injured trauma patient. J Trauma 2009;66:386-392

Kheirabadi BS, Edens JW, Terrazas IB et al: Comparison of new hemostatic granules/powders with currently deployed hemostatic products in a lethal model of extremity arterial hemorrhage in swine. J Trauma 2009;66:316-328

Kragh JF Jr, Walters TJ, Baer DG, Fox CJ, Wade CE, Salinas J, Holcomb JB: Survival with emergency tourniquet use to stop bleeding in major limb trauma. Ann Surg 2009;249:1-7

Lacquiere A, Heard AM: Emergency cricothyroidotomy: training is paramount and oxygenation is the aim. Anaesthesia 2009;64:447-448

Moore FA: Tourniquets: Another adjunct in damage control? Ann Surg 2009;249:8-9

Parker PJ: Pre-Hospital Antibiotic Administration. JR Army Med Corps 2009;154:5-9

Schramko AA, Suojaranta-Ylinen RT, Kuitunen AH, Kukkonen SI, Niemi TT: Rapidly degradable hydroxyethyl starch solutions impair blood coagulation after cardiac surgery: a prospective randomized trial. Anesth Analg 2009;108:30-36

Shipman N, Lessard CS: Pressure applied by the emergency Israeli bandage. Mil Med 2009;174:86-92

Sise MJ, Shackford SR, Sise CB, et al: Early intubation in the management of trauma patients; indications and outcomes in 1,000 consecutive patients. J Trauma 2009;66:32-9

Spinella PC, Perkins JG, Grathwohl KW, Beekley AC, Holcomb JB: Warm fresh whole blood is independently associated with improved survival for patients with combat-related traumatic injuries. J Trauma 2009;66:S69-S76

Stein DM, Dutton RP, Kramer ME, Scalea TM: Reversal of coagulopathy in critically ill patients with traumatic brain injury: recombinant factor VIIa is more cost-effective than plasma. J Trauma 2009;66:63-72

Stevens RL, Rochester AA, Busko J, et al: Needle thoracostomy for tension pneumothorax: failure predicted by chest computed tomography. Prehosp Emerg Care 2009;13:14-17

Talving P, Benfield R, Hadjizachari P, et al: Coagulopathy in severe traumatic brain injury: a prospective study J Trauma 2009;66:55-61

Thistlethwaite R, Ray R, Kessell G: Emergency cricothyroidotomy: equipment in anaesthetic practice. Anaesthesia 2009;64:448-449

Velmahos GC, Tabbara M, Spaniolas K, et al: Self-expanding hemostatic polymer for control of exsanguinating extremity bleeding. J Trauma 2009;66:984-8

Warner KJ, Sharar SR, Copass MK, Bulger EM: Prehospital management of the difficult airway: a prospective cohort study. J Emerg Med 2009;36:257-265

February 2009

Cearnal L: Emergency medicine physicians highlight “Army-wide emergency medical care deficiency” in training of flight medics. Ann Emerg Med 2008;52:522-524

Gerhardt RT, DeLorenzo RA, Olivier J, Holcomb JB, Pfaff JA: Out-of-hospital combat casualty care in the current war in Iraq. Ann Emerg Med 2009;53:169-174

Kragh JF, Walters TJ, Baer, DJ, et al: Survival with emergency tourniquet use to stop bleeding in major limb trauma. Ann Surg 2009;249:1-7

Shipman N, Lessard CS: Pressure applied by the Emergency/Israeli bandage. Milit Med 2009;174:86-92

Suter RE: Forward medical care and the military emergency medicine workforce: too much demand and not enough supply? Ann Emerg Med 2009;53:175-177

Wamer KJ, Cuschieri J, Garland B, et al: The utility of early end-tidal capnography in monitoring ventilation status after severe injury. J Trauma 2009;66:26-31

November 2008

Brodie S, Hodgetts TJ, Ollerton J, et al: Tourniquet use in combat trauma: UK military experience. J R Army Med Corps 2007;153:310-313

Calhoun JH, Murray CK, Manring MM: Multidrug-resistant organisms in military wounds from Iraq and Afghanistan. Clin Orthop Relat Res 2008;466:1356-1362

Englehart MS, Cho SD, Tieu BH, et al: A novel highly porous silica and chitosan-based hemostatic dressing is superior to HemCon and gauze sponges. J Trauma 2008;65:890-892

Hess JR, Brohi K, Dutton RP, et al: The coagulopathy of trauma: a review of mechanisms. J Trauma 2008;65:748-754

Holcomb JB, Wade CE, Michalek JE, et al: Improved plasma and platelet to red blood cell ratios improves outcome in 466 massively transfused civilian trauma patients. Ann Surg 2008;248:in press

Ivascu FA, Howells GA, Junn FS, et al: Predictors of mortality in trauma patients with intracranial hemorrhage on preinjury aspirin or clopidogrel. J Trauma 2008;65:785-788

Kalish J, Burke P, Feldman J, et al: The return of tourniquets. JEMS 2008;33:44-46

Kozek-Langenecker SA, Jungheinrich C, Sauermann W, Van der Linden P: The effects of hydroxyethyl starch 130/0.4 (6%) on blood loss and use of blood products in major surgery: a pooled analysis of randomized clinical trials. Anesth Analg 2008;107:382-390

Lier H, Krep H, Schroeder S, Stuber F: Preconditions of hemostasis in trauma: a review. The influence of acidosis, hypocalcemia, anemia, and hypothermia on functional hemostasis in trauma. J Trauma 2008;65:951-960

Moskowitz DM, Shander A, Javidroozi M, et al: Postoperative blood loss and transfusion associated with use of Hextend in cardiac surgery patients at a blood conservation center. Transfusion 2008;48:768-775

Nichol G, Thomas E, Callaway, et al: Regional variation in out-of-hospital cardiac arrest incidence and outcome. JAMA 2008;300:1423-1431

Rathinam S, Beetman A, Steyn RS: Needle thoracostomy in treatment of a tension pneumothorax. J Trauma 2008;65:964

Sasson C, Hegg AJ, Macy M, et al: Prehospital termination of resuscitation in cases of refractory out-of-hospital cardiac arrest. JAMA 2008;300:1432-1438

Stiver SI, Manley GT: Prehospital management of traumatic brain injury. Neurosurg Focus 2008;25:E5

Tien HC, Jung V, Rizoli SB, Acharya SV, MacDonald JC: An evaluation of Tactical Combat Casualty Care interventions in a combat environment. J Am Coll Surg 2008;207:174-178

Weeks DL, Jahr JS, Lim JC, Butch AW, Driessen B: Does Hextend impair coagulation compared to 6% hetastarch? An ex-vivo thromboelastography study. Am J Ther 2008;15:225-230

August 2008

Adams BD, Cuniowski PA, Muck A, De Lorenzo RA: Registry of emergency airways arriving at combat hospitals. J Trauma 2008;64:1548-1554

Beekley AC: Damage control resuscitation: a sensible approach to the exsanguinating surgical patient. Crit Care Med 2008;36:S267-S274

Beninati W, Meyer MT, Carter TE: The critical care air transport program. Crit Care Med 2008;36:S370-S376

Blackbourne LH: Combat damage control surgery. Crit Care Med 2008;36:S304-S310

Brethauer SA, Chao A, Chambers LW, Green DJ, Brown C, Rhee P, Bohman HR: Invasion vs Insurgency: US Navy/Marine Corps Forward Surgical Care During Operation Iraqi Freedom. Arch Surg. 2008;143(6):564-569.

Dewitt DS, Prough DS: Blast-induced brain injury and post-traumatic hypotension and hypoxemia. J Neurotrauma 2008: E-pub ahead of print 30 April

Doyle GS, Taillac PP: Tourniquets: A review of current use with proposals for expanded prehospital use. Prehospital Emergency Care 2008;12:241-256

Fergusson DA, McIntyre L: The future of clinical trials evaluating blood substitutes. JAMA e-pub 28 April 2008

Kheirabadi BS, Crissey JM, Deguzman R, et al: Effects of synthetic versus natural colloid resuscitation on inducing dilutional coagulopathy and increasing hemorrhage in rabbit. J Trauma 2008;64:1218-1228

Johannigman JA: Maintaining the continuum of en route care. Crit Care Med 2008;36:S377-S382

Kortbeek JB, et al: Advanced Trauma Life Support, 8th Edition, The evidence for change. J Trauma 2008;64:1638-1650

Kozek-Langenecker SA, Jungheinrich C, Sauermann W, Van der Linden P: The effects of hydroxyethyl starch 130/0.4 (6%) on blood loss and use of blood products in major surgery: a pooled analysis of randomized clinical trials. Anesth Anal 2008;107:382-390

Mabry R, McManus JG: Prehospital advances in the management of severe penetrating trauma. Crit Care Med 2008;36:S258-266

Malchow RJ, Black IH: The evolution of pain management in the critically ill trauma patient: emerging concepts from the global war on terrorism. Crit Care Med 2008;36:S346-S347

McIntosh SE, Swanson ER, Barton ED: Cricothyroidotomy in air medical transport. J Trauma 2008;64:1543-1547

Murray CK: Infectious disease complications of combat-related injuries. Crit Care Med 2008;36:S358-S364

Natanson C, Kern SJ, Lurie P, Banks SM, Wolfe SM: Cell-free hemoglobin-based substitutes and risk of myocardial infarction and death. JAMA e-pub 28 April 2008

Niles SE, McLaughlin DF, Perkins JG, Wade CE, Li Y, Spinella PC, Holcomb JB: Increased mortality associated with the early coagulopathy of trauma in combat casualties. J Trauma 2008;64:1459-1463

Perkins JG, Cap AP, Weiss BM, Reid TJ, Bolan CE: Massive transfusion and nonsurgical hemostatic agents. Crit Care Med 2008;36:S325-S329

Richey SL: Tourniquets for the control of traumatic hemorrhage: a review of the literature. World Journal of Emergency Surgery 2006

April 2008

Beekley AC, Sebesta JA, Blackbourne LH, Herbert GS, Kauvar DS, Baer DG, Walters TJ, Mullenix PS, Holcomb JB: Prehospital tourniquet use in Operation Iraqi Freedom: effect on hemorrhage control and outcomes. J Trauma 2008;64:S28-S37

Calhoun JH, Murray CK, Manring MM: Multidrug-resistant organisms in military wounds from Iraq and Afghanistan. Clin Orthop Relat Res 2008; E-pub ahead of print

Cancio LC, Wade CE, West SA, Holcomb JB: Prediction of mortality and of the need for massive transfusion in casualties arriving at combat support hospitals in Iraq. J Trauma 2008;64:S51-S55

Conger NG, Landrum ML, Jenkins DH, et al: Prevention and management of infections associated with combat-related thoracic and abdominal cavity injuries. J Trauma 2008;64:S257-S264

Convertino VA, Ryan KL, Rickards CA, Salinas J, McManus JG, Cooke WH, Holcomb JB: Physiological and medical monitoring for en route care of combat casualties. J Trauma 2008;64:S342-S353

Cordts PR, Brosch LA, Holcomb JB: Now and then: combat casualty care policies for Operation Iraqi Freedom and Operation Enduring Freedom compared with those of Vietnam. J Trauma 2008;64:S14-S20

Dayan L, Zinmann C, Stahl S, Norman D: Complications associated with prolonged tourniquet application on the battlefield. Milit Med 2008;173:63-66

Doyle GS, Taillac PP: Tourniquets – a review of current use with proposals for expanded prehospital use. Prehosp Emerg Care 2008;12:241-256

Hospenthal DR, Murray CK, Anderson RC, et al: Guidelines for the prevention of infection after combat-related injuries. J Trauma 2008;64:S211-S220

Kelly JF, Ritenhour AE, McLaughlin DF, Bagg KA, Apodaca AN, Mallak CT, Pearse L, Lawnick MM, Champion HR, Holcomb JB: Injury severity and causes of death from Operation Iraqi Freedom and Operation Enduring Freedom: 2003-2004 vs 2006. J Trauma 2008;64:S21-S26

Kragh JF, Walters TJ, Baer DG, Fox CJ, Wade CE, Salinas J, Holcomb JB: Practical use of emergency tourniquets to stop bleeding in major limb trauma. J Trauma 2008;64:S38-S50

Lott MN, Fuller JJ, Hancock HA, Singh J, McGwin G, Marcus DM: Vitreal penetration of oral moxifloxacin in humans. Retina 2008;28:473-476

Madigan MC, Kemp CD, Johnson JC, Cotton BA: Secondary abdominal compartment syndrome after severe extremity injury: are early, aggressive fluid resuscitation strategies to blame? J Trauma 2008;64:280-285

Markowitz DM, Shander A, Javidroozi M, Klein JJ, Perelman SI, Nemeth J, Ergin MS: Postoperative blood loss and transfusion associated with use of Hextend in cardiac surgery patients at a blood conservation center. Transfusion 2008;48:768-775

Murray CK, Hsu JR, Solomkin JS, et al: Prevention and management of infections associated with combat-related extremity injuries. J Trauma 2008;64:S239-S251

Murray CK: Epidemiology of infections associated with combat-related injuries in Iraq and Afghanistan. J Trauma 2008;64:S232-S238

Owens BD, Kragh JF, Wenke JC, Macaitis J, Wade CE, Holcomb JB: Combat wounds in Operation Iraqi Freedom and Operation Enduring Freedom. J Trauma 2008;64:295-299

Petersen K, Hayes DK, Blice JP, Hale RG: Prevention and management of infections associated with combat-related head and neck injuries. J Trauma 2008;64:S265-S276

Rhee P, Brown C, Martin M, et al: QuikClot use in trauma for hemorrhage control: case series of 103 documented uses. J Trauma 2008;64:1093-1099

Ritenour AE, Dorlac WC, Fang R, Woods T, Jenkins DH, Flaherty SF, Wade CE, Holcomb JB: Complications after fasciotomy revision and delayed compartment release in combat patients. J Trauma 2008;64:S153-S162

Ryan KL, Batchinsky A, McManus JG, Rickards CA, Convertino VA: Changes in pulse character and mental status are late responses to central hypovolemia. Prehosp Emerg Care 2008;12:192-198

Schwartz D, Amir L, Dichter R, Figenberg Z: The use of a powered device for intraosseous drug and fluid administration in a national EMS: a 4-year experience. J Trauma 2008;64:650-654

Stein DM, Dutton RP, Kramer ME, Handley C, Scalea TM: Recombinant Factor VIIa: decreasing time to intervention in coagulopathic patients with severe traumatic brain injury. J Trauma 2008;64:620-627

Stinger HK, Spinella PC, Perkins JG, et al: The ratio of fibrinogen to red cells transfused affects survival in casualties receiving massive transfusions at an army combat support hospital. J Trauma 2008;64:S79-S85

Varoga D, Drescher W, Lippross S, Pufe T, Schutz R: Nonsteroidal anti-inflammatory drug (NSAID)-related spontaneous compartment syndrome resulting from severe platelet dysfunction. J Traum 2008; E-pub ahead of print

Von Hoff DD, Kuhn JG, Burris HA, Miller LJ: Does intraosseous equal intravenous? A pharmacokinetic study. Am J Emerg Med 2008;26:31-38

Warner KJ, Copass MK, Bulger EM: Paramedic use of needle thoracostomy in the prehospital environment. Prehosp Emerg Care 2008;12:162-168

Wortmann GW, Valadka AB, Moores LE: Prevention and management of infections associated with combat-related central nervous system trauma. J Trauma 2008;64:S252-S256

Yun HC, Branstetter JG, Murray CK: Osteomyelitis in military personnel wounded in Iraq and Afghanistan. J Trauma 2008;64:S163-S168

February 2008

Arnaud F, Tomori T, Saito R, McKeague A, Prusaczyk WK, McKarron RM: Comparative efficacy of granular and bagged formulations of the hemostatic agent QuikClot. J Trauma 2007;63:775-782

Borgman MA, Spinella PC, Perkins JG, Grathwohl KW, Repine T, Beekley AC, Sebesta J, Jenkind D, Wade CE, Holcomb JB: The ratio of blood products transfused in patients receiving massive transfusions at a combat support hospital. J Trauma 2007;63:805-813

Curtis KM, Henriques HF, Fanciullo G, Reynolds CM, Suber F: A fentanyl-based pain management protocol provides early analgesia for adult trauma patients. J Trauma 2007;63:819-826

Guyette FX, Wang H, Cole JS: King airway use by air medical providers. Prehospital Emergency care 2007;11:473-476

Harcke HT, Pearse LA, Levy AD, Getz JM, Robinson SR: Chest wall thickness in military personnel: Implications for needle thoracentesis in tension pneumothorax. Milit Med 2007;172:1260-1263

Kozen BG, Kircher SJ, Henao J, Godinez FS, Johnson AS: An alternative hemostatic dressing: comparison of Celox to HemCon and QuikClot. Acad Emerg Med 2008;15:74-81

McManus J, Hurtado T, Pusateri A, Knoop KJ: A case series describing thermal injury resulting from zeolite use for hemorrhage control in combat operations. Prehospital Emergency Care 2007;11:67-71

Stansbury LG, Lalliss SJ, Branstetter JG, Bagg MR, Holcomb JB: Amputations in U.S. military personnel in the current conflicts in Afghanistan and Iraq. J Orthoo Trauma 2008;22:43-46

Zengerink I, Brink PR, Laupland KB, et al: Needle thoracostomy in the treatment of tension pneumothorax in trauma patients: What size needle? J Trauma 2008;64:111-114

November 2007

Beekley AC, Starnes BW, Sebesta JA: Lessons Learned from modern Military Surgery. Surg Clin N Am 2007;87:157-184

Colyer MH, Weber ED, Weichel ED, Dick JS, Bower KS, Ward TP, Haller JA: delayed intraocular foreign body removal without endophthalmitis during Operations Iraqi Freedom and Enduring Freedom. Ophthalmology 2007;114:1439-1437

Gryth D, Rocksen D, Persson JKE, et al: Severe lung contusion and death after high-velocity behind-armor blunt trauma: relation to protection level. Milit Med 2007;172:1110-1116

Hammett M, Watts D, Hooper T, Pearse L, Naito N: Drowning deaths of U.S. service personnel associated with motor vehicle accidents occurring in Operation Iraqi Freedom and Operation enduring freedom, 2003-2005. Milit Med 2007;172:875-878

Schreiber MA, Perkins J, Kiraly L, Underwood S, Wade C, Holcomb JB: Early predictors of massive transfusion in combat casualties. J Am Coll Surg 2007;205:541-545

Ward KR, Tiba MH, Holbert WH, Blocher CR, Draucker GT, Proffitt EK, Bowlin GL, Ivatury RR, Diegelmann RF: Comparison of a new hemostatic agent to current combat hemostatic agents in a swine model of lethal extremity arterial hemorrhage. J Trauma 2007;63:276-284

August 2007

Eastridge BJ, Jenkins D, Flaherty S, Schiller H, Holcomb JB: Trauma system development in a theater of war: experiences from Operation Iraqi Freedom and Operation Enduring Freedom. J Trauma 2006;61:1366-1372

Gulam H: Care of the Muslim Patient. Journal of Special Operations Medicine 2005;5:39-42

Krag JF, Baer DG, Walters TJ: Extended (16-hour) tourniquet application after combat wounds; a case report and review of the literature. J Orthop Trauma 2007;21:274-278

Lerner EB, O'Connor RE, Schwartz R, et al: Blast-related injuries from terrorism: an international perspective. Prehospital Emergency Care 2007;11:137-153

Perkins JG, Schreiber MA, Wade CE, Holcomb JB: Early versus late recombinant factor VIIa in combat trauma patients requiring massive transfusion. J Trauma 2007;62:1099-1101

Peterson K, Riddle MS, Danko JR, Blazes DL, Hayden R, Tasker SA, Dunne JR: Trauma-related infections in battlefield casualties from Iraq. Ann Surg 2007;245:803-811

Rogatsky GG, Mayevsky A: The life-saving effect of hyperbaric oxygenation during early-phase severe blunt chest injuries. Undersea Hyperb Med 2007;34:75-81